

The  
PSYCHOLOGICAL  
RECORD

Vol. I

NOVEMBER, 1937

No. 23

EFFECT OF BENZEDRINE ON  
ACTIVITY

L. ZIEVE



The Principia Press, Inc.  
Bloomington, Ind.

**EDITOR**

J. R. Kantor  
*Indiana University*

**MANAGING EDITOR**

C. M. Louttit  
*Indiana University*

**DEPARTMENTAL EDITORS****Abnormal**

Edmund S. Conklin,  
*Indiana*

**Child**

Helen Koch,  
*Chicago*

**Clinical**

B. M. Castner,  
*Yale*

**Comparative**

E. A. Culler,  
*Illinois*

**Educational**

J. G. Peatman,  
*C.C.N.Y.*

**Experimental**

B. F. Skinner,  
*Minnesota*

**Medical**

Norman Cameron, M. D.,  
*Johns Hopkins*

**Physiological**

C. F. Scofield,  
*Buffalo*

**Psychometrics**

J. P. Guilford,  
*Nebraska*

**Social**

Norman C. Meier,  
*Iowa*

**Editorial Assistant, J. W. Carter Jr., *Indiana***

---

The Principia Press, Inc., has undertaken the publication of this co-operative journal to afford authors the opportunity of immediate publication at the least possible expense. The present low cost of publication, and possible future reductions, depend entirely upon the number of subscriptions. The subscription price has been made low to induce individuals to subscribe. Under the Articles of Incorporation of the Principia Press no profit can be made on any of its publications. Therefore an increase in the number of subscribers will be reflected in reduced costs to authors and an increase in the number of pages published annually.

**EDITORIAL BOARD.** The above named board of associate editors have agreed to serve in an active capacity for a period of three years, and all manuscripts will be submitted to them.

**MANUSCRIPTS** may be sent to the appropriate associate editor or to Dr. J. R. Kantor. Longer papers (30 or more typewritten pages) in the above mentioned fields are especially desired.

**COSTS.** The cost of publication will be \$2.00 per page of approximately 500 words. For tabular matter or illustrations in excess of 10 per cent of the total length of the paper there will be an excess charge of 75 cents per page. An additional 50 cents will be charged for each paper for the cost of covers.

**REPRINTS.** One hundred copies of each paper will be supplied gratis. Additional copies may be purchased in any quantity at a rate which will be set when the author is informed of the cost of publication.

**ROYALTIES.** Fifty per cent of the net income from the sale of individual copies, or from copies sold as part of back volumes, will be credited as royalties to the author's account. Royalties cannot be paid on income from subscriptions current in the year of publication.

**SUBSCRIPTIONS.** The subscription price is \$4.00 per year for an annual volume of approximately 500 pages. Individual papers will be sold at an advertised price, which will depend upon the size of the article. Foreign subscriptions will be \$4.50.

**CORRESPONDENCE** concerning editorial matters should be addressed to Dr. J. R. Kantor, Indiana University. Business correspondence should be addressed to Dr. C. M. Louttit, Principia Press, Inc., Bloomington, Ind.

## EFFECT OF BENZEDRINE ON ACTIVITY\*

L. ZIEVE

*University of Minnesota*

Casual observation of the immediate after-effects of benzedrine on the human organism indicates a definite hyperkinesis. Greater excitability and a lowered threshold of stimulation are evident. Clinical studies find that benzedrine has a stimulating action on the higher nervous centers; also that it produces a hypertension in the blood vessels. Use is made of its kinetic effect in the treatment of narcoleptic cases (1, 2). Studies of the comparative stimulating effects of various drugs and hormones on the rate of establishment and extinction of a conditioned response illustrate the activating effect of benzedrine (3, 4). The conditioned response was the pressing of a lever. During periodic re-conditioning benzedrine caused an increase of 130 per cent in the rate of responding, and was in this respect unusually effective. It also caused a marked increase in the rate of responding during extinction, being more effective than caffeine.

None of these studies measured spontaneous activity itself. The need for a direct quantitative measurement of the increase in spontaneous activity as a result of ingesting benzedrine prompted this experiment.

The method consisted in running twenty-four three-month-old albino rats in revolving activity drums. These cages have been shown to be reliable (5). The animals were run for three successive 15-day periods, the first and third periods being controls during which no benzedrine was given. An interval of three days was allowed between the second and third periods during which no records were taken though the animals remained in the cages. Each day during the second period 2 cc. of a benzedrine solution were thoroughly mixed with the food. Addition of the drug was made separately to the individual food cups to assure accurate dispensation. The dose amounted to 2 mgm. of benzedrine per kilo of body weight. This amount was chosen as a medium dose for animals weighing about

\*Recommended for publication by Dr. B. F. Skinner, September 25, 1937.

250 gms., after preliminary experimentation with doses varying from .5 mgm. to 40 mgm. per kilo of body weight.<sup>1</sup> The amount of food eaten by the animals was held constant from day to day.

Means were calculated in terms of revolutions of an 8-in. revolving cage for the first control,  $C_1$ , the test run, T, and the second control,  $C_2$ . The results are summarized in Table I.

TABLE I  
REVOLUTIONS OF CAGES IN CONTROL ( $C_1$ ,  $C_2$ ) AND  
BENZEDRINE (T) PERIODS

Period	Mean	$\sigma$	$D/\sigma_D$	$\frac{\sigma}{\text{Mean}}$
$C_1$	62,500	51,900	$C_1:T: 3.54$	83.04%
T	135,000	85,950	$C_2:T: 3.05$	63.67%
$C_2$	70,840	56,840	$C_1:C_2: 0.53$	80.24%

It will be noted that between the first control and the test run there is a critical ratio of 3.54, between the second control and the test run a critical ratio of 3.05, and between the first and second controls a critical ratio of 0.53. The great differences between  $C_1$  and the test run and between  $C_2$  and the test run leave little room for doubt as to the hyperkinetic effect of benzedrine. An increase of more than 100 per cent over the average control activity is produced. These differences are especially significant in view of the fact that only 24 animals were used.

The ratio of 0.53 between  $C_1$  and  $C_2$  represents an important isolating control showing that the increase in activity during the test run was not the result of prolonged stay in the activity cages.

A difference of approximately 20 per cent between the coefficients of variation for the two control periods and the test period brings out a further result; namely, that benzedrine tends to reduce relative variability. The decrease in relative variability resulted from a proportionately greater increase in activity of the less active animals. In other words, benzedrine seems to have the greatest activating effect on the least active animals.

Taking the activity increase as a measure of the physiological effects of benzedrine, one can get some evidence as to the tolerance built up to the drug by dividing the test run, which consisted of records for fifteen consecutive days, into 3 equal sub-periods and comparing the records for each sub-period. Means computed for these sub-periods are given in Table II.

<sup>1</sup> The experimentation on varying doses was largely the work of Mr. H. Crook.

TABLE II  
REVOLUTIONS OF CAGES DURING SUCCESSIVE PARTS OF  
THE BENZEDRINE PERIOD

Five-day Sub-periods	Mean	$\sigma$	$\sigma$
			Mean
First	36,667	23,570	64.28%
Second	42,917	29,577	68.92%
Third	48,333	35,668	73.80%

It would be expected that with the development of tolerance to the drug there would be a decrease in activity unless the dosage is very near the depressant stage, which is unlikely. Comparison of the records in Table II of the three sub-periods indicates no decrease, but rather a slight though insignificant increase. It is, therefore, probable that the animal does not build up a tolerance to the drug.

In the light of the following data, there seems to be little likelihood that an effect of the dose beyond 24 hours needs to be considered here. The mean number of revolutions for the last day of the benzedrine period was 11,250, for the following day with no benzedrine 3,167, and for the first day of the second control period 3,917.<sup>2</sup> The respective sigmas were 8,319, 2,779, and 3,226. Such a sudden decrease, notwithstanding the large standard deviations, stands in contradistinction to the fairly gradually descending curve that one would expect had there been a cumulative effect.

In Table II are listed coefficients of variation that appear to have uniformly increasing values. This seemingly significant orderliness is probably an artifact resulting from the arbitrary method of dividing the test period into sub-periods. When the test period was broken up into three-day intervals so that there were five sub-periods instead of three, the coefficients of variation showed a chance distribution. Thus the coefficients for the first, second, third, fourth, and fifth sub-periods were respectively 71.50, 61.82, 66.62, 72.63, and 69.81 per cent.

Incidental observations on the effects of large doses of benzedrine (40 mgm. per kiloweight) on activity indicate that it has a depressant effect. It was difficult to get very accurate records because interfering visceral inhibitory and sudorific effects were marked.

<sup>2</sup> The second control began on the fourth day following the test period. During the intervening time the animals remained in the cages.

## SUMMARY

Records of the spontaneous activity of animals were accumulated over a period of 15 days during which a medium dose of benzedrine was ingested daily. These records were compared with similar records of two control periods of the same length, one preceding and one following the benzedrine test period. The significant differences between the control and test periods definitely illustrate the activating properties of benzedrine. A difference of approximately 20 per cent between the relative variability of the control periods and the test period indicates that benzedrine tends to reduce relative variability. Evidence obtained indicated the probability that no tolerance was built up to the drug.

## REFERENCES

1. Prinzmetal, M., and Bloomberg, W. The use of benzedrine for the treatment of narcolepsy. *J. A. M. A.*, 1935, 105; 2051-2054.
2. Peoples, S. A., and Guttman, E. Hypertension produced with benzedrine. *Lancet*, 1936, 20; 1107-1109.
3. Wentink, E. Effect of certain drugs and hormones on conditioning. *J. Exp. Psych.* To be published.
4. Skinner, B. F., and Heron, W. T. Effects of caffeine and benzedrine upon conditioning and extinction. *Psych. Record*. 1937, 1; 339-346.
5. Shirley, M. Studies in activity. I. Consistency of the revolving drum method of measuring the activity of the rat. *J. Comp. Psych.*, 1928, 8; 23-38.

